

Modifiable and Non-modifiable Factors Associated with Low Awareness of Hypertension Treatment in Indonesia: A Cross-Sectional Population-Based National Survey



QISTY A. KHOIRY

SOFA D. ALFIAN

RIZKY ABDULAH

*Author affiliations can be found in the back matter of this article

Ubiquity press

ABSTRACT

Introduction: Low awareness about hypertension treatment is recognized as a significant cause of treatment failure. Therefore, identifying its underlying factors is essential for developing effective intervention strategies. This study aims to identify the modifiable and non-modifiable factors associated with low awareness about hypertension treatment.

Method: This national, cross-sectional, population-based survey used publicly available data from the Indonesian Family Life Survey (IFLS-5) for 2014 among respondents with hypertension aged ≥ 15 years. Depression and insomnia, as modifiable factors, were assessed using the Centre for Epidemiologic Studies—Depression (CES-D) and the Patient-Recorded Outcomes Measurement Information System (PROMIS) questionnaire, respectively. Non-modifiable factors, such as sociodemographic information, were obtained from self-reported data. Logistic regression analysis was used to assess the association between these factors and low awareness about hypertension treatment. Odds ratios (ORs) with 95% confidence intervals (CIs) were reported.

Result: The study recruited 7,920 respondents, the majority of whom were female (53.8%) and aged < 60 years (71.1%). The prevalence of low awareness of hypertension treatment was 87.1% (51.8% in women and 48.2% in men). Being an elderly (OR: 1.60, 95%CI 1.36–1.88), being irregularly blood pressure control (OR: 4.40, 95% CI 3.78–5.13), having depressive symptoms (OR: 1.35, 95% CI 1.12–1.62), having insomnia (OR: 1.31, 95% CI 1.11–1.53), and having low satisfaction with health care (OR: 1.28, 95% CI 1.08–1.51) were associated with low awareness of hypertension treatment. Surprisingly, respondents with strong religiosity (OR: 1.62; 95% CI 1.25–2.09) were more likely to display low awareness of hypertension treatment.

Conclusion: The main factors associated with low awareness of hypertension treatment are modifiable. Thus, health care professionals should integrate more patient-specific factors when designing tailored interventions.

CORRESPONDING AUTHOR:

Sofa D. Alfian

Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy, Universitas Padjadjaran, Jatinangor, Indonesia; Center of Excellence in Higher Education for Pharmaceutical Care Innovation, Universitas Padjadjaran, Jatinangor, Indonesia

sofa.alfian@unpad.ac.id

KEYWORDS:

Awareness; Hypertension; Medication; Treatment

TO CITE THIS ARTICLE:

Khoiry QA, Alfian SD, Abdulah R. Modifiable and Non-modifiable Factors Associated with Low Awareness of Hypertension Treatment in Indonesia: A Cross-Sectional Population-Based National Survey. *Global Heart*. 2022; 17(1): 56. DOI: <https://doi.org/10.5334/gh.1143>

INTRODUCTION

Hypertension is an increasing part of the global burden of disease, which is more predominant in low- and middle-income countries (LMICs) [1]. The current global trends in hypertension has increased from 650 million to 1.28 billion over the last three decades [2]. A population-based study on more than 90 countries revealed that the burden of hypertension has increased by 7.7% in LMICs but decreased by 2.6% in high-income countries (HICs) [3]. Globally, 8.5 million deaths were associated with uncontrolled high blood pressure due to unawareness of this condition [4]. In fact, awareness of hypertension treatment is reportedly lower in LMICs compared with other countries [5]. In 2017, Indonesia reported 91.3 million people with hypertension, which was responsible for 35% of all deaths [6, 7].

Uncontrolled blood pressure is predominantly frequent in LMICs, where health awareness is the weakest and mortality rates due to cardiovascular disease are the highest [8]. According to Indonesia Basic Health Research 2018, only 37% of Indonesians with hypertension currently have obtained control of their blood pressures, whereas 45.6% exhibited low adherence to hypertension treatment [7]. Low awareness of medication is reported to be associated with medication non-adherence [9, 10], which may lead to poorly controlled hypertension, high risks of complication, and increased health care costs [11–13]. Therefore, low awareness of hypertension treatment should be addressed due to the lack of symptoms of the disease.

Previous studies have reported several factors that may be associated with awareness of hypertension treatment [14–17]. However, such studies have mainly explored non-modifiable factors. A previous study conducted in LMICs revealed that insurance status and income were significantly associated with awareness of treatment of hypertension [18]. Moreover, low levels of education were associated with low rates of awareness about hypertension treatment in LMICs but not in HICs [5]. However, the association of modifiable factors, such as satisfaction with health care, blood pressure control status, depressive symptoms, and insomnia, to awareness of hypertension treatment in Indonesia remains unclear.

The Indonesian Family Life Survey (IFLS) is a longitudinal socioeconomic and health survey in Indonesia, which represents approximately 83% of the Indonesian population [19]. Several studies that utilized IFLS-4 or IFLS-5 demonstrated the association of non-modifiable factors, such as socioeconomic status [20] and psychosocial risk [17], to hypertension in Indonesia. However, a clear evidence of which factors or foci are required to improve medication awareness of hypertension treatment in Indonesia remains lacking. Thus, identifying its underlying factors is essential to developing effective strategies for intervention. This study aims to identify modifiable and non-modifiable factors associated with low awareness of hypertension treatment in Indonesia.

METHODS

STUDY DESIGN

This study used the national longitudinal and cross-sectional data of the fifth wave of the IFLS-5, which were collected from 2014 to 2015 and were publicly available since 2016. IFLS-5 used a multistage stratified sampling design that represents 83% of the Indonesian population conducted in 13 provinces, which are four provinces on Sumatra (North Sumatra, West Sumatra, South Sumatra, and Lampung), all five of the Javanese provinces (DKI Jakarta, West Java, Central Java, DI Yogyakarta, and East Java), and four provinces covering the remaining major island groups (Bali, West Nusa Tenggara, South Kalimantan, and South Sulawesi) with a response rate of more than 90% [19]. The IFLS collected extensive measures of basic sociodemographic, certain economic characteristics of all household members, and health status, including self-reported measures of general health status, symptoms, pain, and biomarker measurements of health. The ethical review boards have approved the IFLS of Research and Development (RAND) Corporation and Gadjah Mada University. Written informed consent was obtained from all respondents before data collection [19].

STUDY POPULATION

The total number of IFLS-5 subjects was 34,464 (aged 0 to >80 years). Data were obtained from subjects who were at least 15 years of age at the completion of the survey. Subjects with available data on blood pressure, depressive symptoms, and sleeping quality were included. Subjects with missing values for outcome variables were excluded.

Subjects with hypertension were defined as having a blood pressure $\geq 140/90$ mm Hg [21]. Blood pressure was measured three times at home by specially trained nurses using an Omron sphygmomanometer (HEM-7203) as the subjects were seated. The first measurement was taken at the beginning of the interview with two subsequent measures taken during the interview [19]. The average of the three measurements was used for analysis.

Awareness of treatment was defined by responses to the following question: *In order to manage your hypertension, are you currently taking prescribed medication weekly?* Those who answered **yes** were considered to have a high awareness of hypertension treatment, whereas the opposite is true for those who responded with **no**. Awareness of hypertension treatment refers to those who were aware of their condition, aware they need to have treatment, and receiving treatment for their condition [22, 23]. It is relatively different from adherence, which was defined as the process by which patients take their medications as prescribed, composed of initiation (when the first dose is started), implementation (the actual dose of the patient from the start to the last dose), and discontinuation (end of treatment) [24].

POTENTIAL FACTORS ASSOCIATED WITH LOW AWARENESS OF HYPERTENSION TREATMENT

The potential factors associated with low awareness of hypertension treatment were classified as modifiable and non-modifiable factors. Depressive symptoms, having insomnia, blood pressure control status, happiness status, having visited health care services, and satisfaction with health care were categorized into modifiable factors. Alternatively, non-modifiable factors consisted of gender, age, marital status, level of education, residency, coverage of health insurance, economic status, and religiosity.

A self-reported Center for Epidemiologic Studies Depression (CES-D) scale was used to measure depressive symptoms [19]. CES-D contains 10 questions to assess the feelings of subjects, which are highly correlated with depressive symptoms [25]. Eight questions were focused on negative symptoms (e.g., *I was bothered by things that usually do not bother me*), whereas the two other questions assessed the positive symptoms of depression (e.g., *I felt hopeful about the future*). The subjects indicated the frequency that each item applied to them in the past week using a four-point Likert-type scale (0 = rarely or none of the time; 1 = some or little of the time; 2 = moderately or much of the time; 3 = most or almost all the time). The final score is calculated by summing all items after reversing the positive mood items. A subject with a total score equal to or above 10 is considered to have symptoms of depression [26]. The CES-D questionnaire was translated into Indonesian (forward translation), where two independently hired translators re-translated it into English (back translation) [19].

Insomnia was assessed using 10 questions from PROMIS [19]. Five items each were used to assess sleep quality and sleep impairment in the past week [27]. Each item was rated using a five-point Likert-type scale (0 = never/not at all; 1 = a little bit; 2 = somewhat; 3 = quite a bit; 4 = always/very much). Insomnia was defined as total scores of ≥ 21 –40 [28]. The PROMIS questionnaire was translated into Indonesian (forward translation), where two independently hired translators re-translated it into English (back translation) [19].

Blood pressure control status was categorized as regular and irregular dependent on the subject's response to the question, *How regularly do you have your blood pressure checked?* Happiness status was assessed using the question, *Taken together, how you would say things are these days? Would you say you are very happy, happy, unhappy, or very unhappy?* Those who answered *very happy* or *happy* were categorized as happy, whereas those who answered *unhappy* or *very unhappy* were categorized as unhappy [29]. Satisfaction with health care was assessed using the question, *Concerning your health care, which of the following is true: it is less than adequate for my needs; it is just adequate for my needs; it is more than adequate for my needs?* Low satisfaction with health care is equivalent to the response *it is just less than adequate for my needs*, whereas high satisfaction with health care is indicated by *it is just adequate for my needs* or *it is more than adequate for my needs* [29].

Sociodemographic factors were gender (men, women), age at the completion of the survey (less than 60 years for the non-elderly and more than 60 years for the elderly), marital status (currently unmarried and currently married), level of education (no education, elementary school, junior high school, senior high school, and university), residency (rural and urban),

coverage of health insurance (no and yes), economic status (poor and non-poor), and religiosity (religious, non-religious). The total income of the family was used to measure economic status over the past 12 months in rupiah divided by family size (per capita income). Capita income was categorized per quintile. Religiosity was assessed using the question *How religious are you?* Those who answered *very religious* or *religious* were categorized as religious, whereas those who answered *somewhat religious* or *not religious* were considered as non-religious [29]. The subject was asked whether they visited any health care services one month prior to the survey to assess health care visit status (yes and no).

To determine whether subjects have comorbid disease status, subjects were asked the question, *Has a doctor/ paramedic/nurse/midwife ever told you that you had the following chronic conditions of diseases?* with response options: hypertension, diabetes, tuberculosis, asthma and other chronic lung diseases, cardiac diseases(heart attack/coronary heart disease/angina or other heart diseases), liver diseases, stroke, cancer or malignancies, arthritis/rheumatism, uric acid/gout, depression, vision and hearing abnormalities [29]. Subjects who answered only hypertension were categorized as having no comorbidity, while subjects who answered hypertension and one to three other chronic disease were then categorized as having one to three comorbidities, and subjects who had more than four comorbidities were categorized as having more than four comorbidities.

STATISTICAL ANALYSIS

Descriptive statistics were used to summarize the characteristics of the subjects. Awareness of hypertension treatment was estimated for each age group and gender. A chi-square test was performed to assess the univariate association between the characteristics of the subjects and the outcome. We conducted complete case analyses because the missing data were small. The potential factors found associated with the outcome at a significance level of $p < 0.25$ in the univariate analyses were included in the initial multivariate model. Multivariate logistic regression was performed to obtain the odds ratio (OR) with a 95% confidence interval (95% CI) with manual backward elimination. The p -values were set at 0.05 for factors included in the final model. Subgroup analyses by number of comorbidities was performed in no comorbid, having one to three comorbidities, and having more than four comorbidities groups. The goodness-of-fit statistic was assessed using the Hosmer-Lemeshow test. Pseudo R-squared value, which is the default value reported by Stata, is obtained as an equivalent to the R-squared reported in the linear regression. R-squared is a number that ranges from 0 to 1, which indicates the extent to which the combination of independent variables simultaneously influences the value of the dependent variable [30]. All statistical analyses were performed using Stata software version 14.0 for Windows.

RESULTS

A total sample of 7,920 subjects without missing data on awareness of hypertension treatment was included in this study. The majority were female (53.8%) and aged <60 years (71.1%; Table 1). A high proportion of subjects displayed poor blood pressure control (80.5%), although most of them (90.8%) visited health care facilities in the last month. Approximately half of subjects were covered by health insurance (50.4%), resided in rural areas (57.6%), and graduated from elementary school (41.2%). Out of the total participants, 20.4% experienced depression, whereas 46.3% have insomnia.

CHARACTERISTIC	NUMBER	%
Awareness of Hypertension Treatment		
Low Awareness	6,895	87.1
High Awareness	1,025	12.9
Comorbidity Status		
No Comorbid	5,134	67.1
1-3 Comorbidities	2,441	31.8
≥4 Comorbidities	81	1.1
Missing	264	—
Gender		
Women	4,263	53.8
Men	3,655	46.2
Missing	2	—

Table 1 Baseline Characteristic of Study Population (N = 7,920).

(Contd.)

CHARACTERISTIC	NUMBER	%
Age		
Non-elderly [<60 years old]	5,627	71.1
Elderly [≥60 years old]	2,291	28.9
Missing	2	—
Education Level		
No Education	1,021	13.3
Elementary School	3,173	41.2
Junior High School	1,099	14.3
Senior High School	200	2.6
University	2,208	28.7
Missing	219	—
Marital Status		
Not Currently Married	2,138	27.0
Currently Married	5,780	73.0
Missing	2	—
Blood Pressure Control Status		
Irregularly	4971	80.5
Regularly	1,201	19.5
Missing	1,748	—
Economic Status		
Quintile 1	1,605	20.3
Quintile 2	1,661	21.0
Quintile 3	1,498	18.9
Quintile 4	1,595	20.1
Quintile 5	1559	19.7
Missing	2	—
Health Care Satisfaction		
Low	1,837	25.3
High	5,411	74.7
Missing	672	—
Happiness Status		
Unhappy	857	11.8
Happy	6,391	88.2
Missing	672	—
Insomnia		
Yes	3,353	46.3
No	3,895	53.7
Missing	672	—
Depressive Symptoms Status		
Not Depressed	5,768	79.6
Depressed	1,480	20.4
Missing	672	—
Residency		
Rural	4,460	57.6
Urban	3,289	42.4
Missing	171	—
Health Care Visit Status		
No	1,548	90.8
Yes	156	9.2
Missing	6,216	—
Coverage of Health Insurance		
No	2,105	49.6
Yes	2,141	50.4
Missing	3,674	—
Religiosity		
Not Religious	1,248	17.2
Religious	6,002	82.8
Missing	670	—

The prevalence of low awareness of hypertension treatment was 87.1% (51.8% in women and 48.2% in men). Level of education, marital status, economic status, and happiness status were negatively associated with low awareness of hypertension treatment (Table 2). Gender, age, blood pressure control status, satisfaction with health care, insomnia, depressive symptoms status, residency, health care visit status, coverage of health insurance, and religiosity were selected as potential factors associated with low awareness of hypertension treatment based on univariate analyses.

CHARACTERISTIC	LOW AWARENESS OF HYPERTENSION TREATMENT		HIGH AWARENESS OF HYPERTENSION TREATMENT		P-VALUE
	N	%	N	%	
Gender					0.000*
Women	3,568	51.8	695	67.9	
Men	3,326	48.2	329	32.1	
Comorbidity Status					0.000*
No Comorbid	4,865	70.5	269	35.3	
1–3 Comorbidities	1,977	28.7	464	61	
≥4 Comorbidities	53	0.8	28	3.7	
Age					0.000*
Non-elderly [<60 years old]	5,007	72.6	620	60.5	
Elderly [≥60 years old]	1,887	27.4	404	39.5	
Education Level					0.368
No Education	876	13.1	145	14.5	
Elementary School	2,761	41.2	412	41.3	
Junior High School	960	14.3	139	13.9	
Senior High School	182	2.7	18	1.8	
University	1,924	28.7	284	28.5	
Marital Status					0.910
Not Currently Married	1,860	26.9	278	27.2	
Currently Married	5,034	73.1	746	72.8	
Economic Status					0.251
Quintile 1	1,396	20.3	209	20.4	
Quintile 2	1,462	21.2	199	19.4	
Quintile 3	1,309	19.0	189	18.5	
Quintile 4	1,395	20.2	200	19.5	
Quintile 5	1,332	19.3	227	22.2	
Residency					0.000*
Rural	2,924	42.4	660	64.4	
Urban	3,971	57.6	365	35.6	
Health Care Visit Status					0.000*
No	1,046	88.6	502	96.0	
Yes	135	11.4	21	4.0	
Coverage of Health Insurance					0.000*
No	1,892	51.0	320	60.0	
Yes	1,821	49.0	213	40.0	
Religiosity					0.000*
Not Religious	1,139	18.0	107	11.7	
Religious	5,191	82	811	88.3	
Blood Pressure Control Status					0.000*
Irregularly	5,328	84.2	514	56.1	
Regularly	1,001	15.8	403	43.9	

Table 2 Univariate Associations with Awareness of Hypertension Treatment.

* Included in initial multivariate model.

(Contd.)

CHARACTERISTIC	LOW AWARENESS OF HYPERTENSION TREATMENT		HIGH AWARENESS OF HYPERTENSION TREATMENT		P-VALUE
	N	%	N	%	
Health Care Satisfaction					0.000*
Low	1,553	24.5	284	30.9	
High	4,777	75.5	634	69.1	
Happiness Status					0.210
Unhappy	737	11.6	120	13.1	
Happy	5,593	88.4	798	86.9	
Insomnia					0.000*
Yes	2,858	45.2	423	46.1	
No	3,472	54.8	495	53.9	
Depressive Symptoms Status					0.000*
Not Depressed	5,091	80.4	677	73.7	
Depressed	1,239	19.6	241	26.3	

In the multivariate model, the elderly (OR: 1.60; 95% CI: 1.36–1.88) and subjects with low satisfaction with health care (OR: 1.28; 95% CI: 1.08–1.51) were associated with low awareness of hypertension treatment. Interestingly, subjects with high religiosity were likely to exhibit low awareness of hypertension treatment (OR: 1.55; 95% CI: 1.24–1.94). Modifiable factors associated with low awareness of hypertension treatment were having depressed symptoms (OR: 1.35; 95% CI: 1.12–1.62), having insomnia (OR: 1.31; 95% CI: 1.11–1.53), and blood pressure control status (OR: 4.40; 95% CI: 3.78–5.13). Irregular blood pressure control exhibited the highest odds of low awareness of hypertension treatment (Table 3). The goodness-of-fit p-value of the model was 0.271 with a pseudo-R-squared value of 8.74%. Overall, modifiable and non-modifiable factors that were associated with low awareness of hypertension treatment were relatively similar in subgroup analysis, except that insomnia and depressive symptom status were not seen as risk factors for low awareness of hypertension treatment among subjects with more than four comorbidities (Table 3).

Table 3 Association between Modifiable and Non-modifiable Factors with Low Awareness of Hypertension Treatment with Subgroup Analysis Result.

^a Goodness-of-fit p-value: 0.271; pseudo-R-squared: 8.74%.

^b Final multivariate model.

* Not significant.

CHARACTERISTIC	OVERALL ^a		SUBGROUP ANALYSIS: NO COMORBIDITIES		SUBGROUP ANALYSIS: 1–3 COMORBIDITIES		SUBGROUP ANALYSIS: ≥4 COMORBIDITIES	
	OR [95% CI] ^b	P-VALUE	OR [95% CI]	P-VALUE	OR [95% CI]	P-VALUE	OR [95% CI]	P-VALUE
Age								
Non-elderly (<60 years old)	Reference		Reference		Reference		Reference	
Elderly (≥60 years old)	1.60 [1.36–1.88]	0.000	1.72 [1.30–2.29]	0.000	1.82 [1.42–2.32]	0.000	1.34 [0.39–4.61]	0.641*
Blood Pressure Control Status								
Regularly	Reference		Reference		Reference		Reference	
Irregularly	4.40 [3.78–5.13]	0.000	3.47 [2.63–4.60]	0.000	4.48 [3.54–5.69]	0.000	2.14 [0.64–7.16]	0.219*
Depressive Symptoms Status								
Not Depressed	Reference		Reference		Reference		Reference	
Depressed	1.35 [1.12–1.62]	0.002	1.04 [0.73–1.48]	0.821*	1.37 [1.04–1.82]	0.028	0.50 [0.10–2.47]	0.398*
Insomnia								
No	Reference		Reference		Reference		Reference	
Yes	1.31 [1.11–1.53]	0.001	1.46 [1.02–2.09]	0.041	1.19 [0.86–1.66]	0.293*	1.09 [0.21–5.74]	0.916*
Health Care Satisfaction								
High	Reference		Reference		Reference		Reference	
Low	1.28 [1.08–1.51]	0.003	1.27 [0.94–1.71]	0.114*	1.14 [0.88–1.47]	0.324*	0.60 [0.16–2.21]	0.442*
Religiosity								
Not Religious	Reference		Reference		Reference		Reference	
Religious	1.55 [1.24–1.94]	0.000	1.75 [1.14–2.68]	0.011	1.56 [1.09–2.22]	0.016	2.29 [0.38–13.70]	0.036*

Among the 7,920 respondents, 87.1% reported low awareness of hypertension treatment. Age and religiosity are non-modifiable factors associated with low awareness of hypertension treatment. Meanwhile, the modifiable factors are blood pressure control status, satisfaction with health care, depressive symptoms, and insomnia. Medication awareness is critical in providing accurate information about the treatment undertaken to improve medication adherence [31]. Patients with low awareness of treatment tend to be non-adherent to treatment for chronic diseases [32]. Medication awareness is one of the five broad categories of treatment-related problems in a patient perspective that have potential implications for therapeutic success as they create barriers to medication adherence [33]. However, low adherence to treatment in chronic disease is associated with poor disease control, increased costs, increased hospitalization rates, decreased quality of life, and increased mortality [34–37].

We observed that the elderly subjects were associated with low awareness of hypertension treatment, which can be partially explained by several possible barriers, such as frailty, multiple comorbidities, polypharmacy, and cognitive impairment [38]. Surprisingly, religiosity indicated a significant association with low awareness of hypertension treatment. A possible explanation may be that people foresee and surrender healing outcomes from God. Thus, they would risk not being aware of taking medications [39, 40]. In addition, previous studies reported that religiosity could increase the likelihood of using complementary and alternative medical (CAM) practices, such as herbal medicines [41, 42]. The high prevalence of CAM among patients in Ethiopia with hypertension is associated with low awareness of health care, which potentially leads to ineffective hypertension management [43]. Another finding from a previous study reported decreased adherence to prescribed medication due to CAM use in patients with type 2 diabetes [44].

We further observed that blood pressure control status, satisfaction with health care, depressive symptoms, and insomnia are associated with low hypertension awareness. Subjects who irregularly control their blood pressure have a risk of low awareness of hypertension treatment. A possible explanation for this finding may be that the inadequate understanding of therapeutic goals influences their awareness of hypertension treatment [45]. Affordability and unreliable devices are the two main reasons why patients do not check their blood pressure regularly [46]. Individual medical education is the most critical aspect of blood pressure regulation [47]. The fundamental method for hypertension control is to make people more aware of it, treat it, and maintain it under control [48]. The problem of achievement of target blood pressure consists of patient-related barrier, physician-related barrier, and health care system-related barrier, so that a team-based approach to blood pressure control that involves nurses, pharmacists, and physician assistants should be emphasized [49]. People with high blood pressure can benefit from antihypertensive medication, while complications and aging make it less probable to achieve the desired blood pressure [47].

We also observed that patients with low satisfaction with health care were likely to display low awareness of hypertension treatment. Previous studies revealed that satisfaction with health care was associated with adherence to hypertension treatment [50, 51]. Subjects with better satisfaction with health care may increase their awareness of hypertension treatment. Furthermore, having a depressive status was associated with low awareness of hypertension treatment, which can be partially explained by the deterioration of high-level cognitive functions, specifically attention, awareness, and memory, among subjects with depressive symptoms [52]. At the same time, having depressive disorders has been associated with hypertension [53–55]. A systematic review and meta-analysis demonstrated that the risk for cardiovascular diseases in subjects with depressive symptoms is approximately two times higher than those in non-depressed people [56]. Similar results were observed in subjects with insomnia. A possible explanation is that the severe decrease in sleep hours is related to cognitive function, which leads to the lack of awareness of treatment. Prior studies have reported a significant relationship between sleep duration and cognitive function [57, 58]. This finding indicates the need to screen for sleep impairment and to maintain regular sleep patterns.

Hypertension frequently coexists with diabetes, hyperlipidemia, or other metabolic syndromes [59]. Patients with these comorbidities are more likely to require combination therapy, which needs more medications. The result of subgroup analysis showed that both subgroups had

relatively similar modifiable and non-modifiable factors that were associated with low awareness of hypertension treatment. Therefore, health care providers need to pay more attention to patients with low awareness regardless of their number of comorbidities.

In summary, the majority of sociodemographics as non-modifiable factors were not associated with low awareness of hypertension treatment. Sociodemographics such as marital status may be extremely general for predicting the medication awareness of subjects. No association was found between gender and low awareness of hypertension, as reported in a previous study [60]. In general, gender is seemingly a less meaningful factor associated with low awareness of hypertension treatment. Moreover, the level of education is not associated with awareness, which is similar to the findings of previous studies [61, 62]. Thus, health literacy may be more important than level of education [62]. By contrast, another study indicated that low levels of education were associated with low awareness of hypertension in LMICs [5]. We observed that residency was not associated with low awareness of hypertension treatment. This finding is in contrast with that of a previous study [63], which found an association between people living in urban areas and high awareness of hypertension treatment. Population in urban areas has exhibited better treatment-seeking behavior and has easy access to health care [63]. Notably, the current study observed no association between visits to health care facilities with low awareness of hypertension treatment. In contrast, a study using IFLS-4 indicated that the number of visits to any health care facility helped to increase awareness about hypertension [64]. Furthermore, no association was found between happiness status and low awareness of hypertension treatment. This result is consistent with data obtained in a cohort study with a 31-year follow-up in London, which revealed the lack of robust associations between life satisfaction and incident hypertension [65]. One unexpected finding is that health insurance coverage was not associated with low awareness of hypertension treatment. However, one could expect that the limited subjects had health coverage in 2014 since the Indonesian National Health Insurance System began in 2014. Moreover, other studies reported on the positive effect of health insurance coverage due to the reduced cost of receiving health care services [66, 67].

The current finding suggests that the majority of non-modifiable factors may be insufficient to address the low awareness of hypertension treatment. Thus, health care professionals should consider modifiable factors in designing tailored and personalized interventions by implementing open communication to educate and to understand patients' views regarding patients' and family members' involvement in the treatment to improve patients' health outcomes [68]. Education and counseling are effective in terms of behavioral changes in patients to clarify misperceptions and improve the awareness of hypertension treatment [69]. Furthermore, health care professionals should regularly check blood pressure and consider patient satisfaction by providing correct treatment and creating a respectful and caring relationship with patients by sharing the decision-making process [70]. The current findings may serve as reference for health professionals focusing on factors related to low awareness of hypertension treatment that should be addressed (Figure 1).

The strength of this study is that we used IFLS data, which represents 83% of the Indonesian population with a small attrition rate (6%). However, we acknowledge certain limitations related to methodological issues. First, the study was unable to provide causal inferences regarding the association between modifiable and non-modifiable factors related to low awareness of hypertension treatment due to the cross-sectional design. Second, we used complete case analysis, which may reduce statistical power due to the sample size, which can produce biased estimates, leading to overestimating or underestimating conclusions. Third, in this analysis, we included only subjects with a blood pressure of more than 140/90 mmHg, which may lead to overestimating low awareness of hypertension treatment since subjects with controlled blood pressure (blood pressure <140/90 mmHg) are not included. Fourth, the overall association of our model was relatively low, which indicated the presence of other unmeasured factors that may influence the low awareness of hypertension treatment, such as education about hypertension, origins of subjects, local tradition, number of medication, medication beliefs [71], comorbid diseases [72], or healthy lifestyle [73].




Figure 1 Problems, Challenges, and Solutions to Increase Low awareness of Hypertension Treatment in Indonesia.

CONCLUSIONS

The main factors associated with low awareness of hypertension treatment are modifiable. Thus, health care professionals should integrate more patient-specific factors when designing tailored interventions.

FUNDING INFORMATION

This study was supported by grant-in-aid from Universitas Padjadjaran. QAK is supported by a postgraduate scholarship (PMDSU Scholarship) from the Indonesian Ministry of Education, Culture, Research, and Technology.

COMPETING INTERESTS

The authors have no competing interests to declare.

AUTHOR CONTRIBUTIONS

QAK wrote the first draft of this manuscript. SDA and RA participated in the design of the study. QAK and SDA participated in data analysis and interpretation. QAK, SDA, and RA contributed to the revision of the manuscript. All authors approved the final manuscript.

Qisty A. Khoiry  orcid.org/0000-0002-4920-2434

Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy, Universitas Padjadjaran, Jatinangor, Indonesia

Sofa D. Alfian  orcid.org/0000-0001-5419-8938

Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy, Universitas Padjadjaran, Jatinangor, Indonesia; Center of Excellence in Higher Education for Pharmaceutical Care Innovation, Universitas Padjadjaran, Jatinangor, Indonesia

Rizky Abdulah  orcid.org/0000-0002-8779-6421

Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy, Universitas Padjadjaran, Jatinangor, Indonesia; Center of Excellence in Higher Education for Pharmaceutical Care Innovation, Universitas Padjadjaran, Jatinangor, Indonesia

REFERENCES

- Zhou B, Perel P, Mensah GA, Ezzati M.** Global epidemiology, health burden and effective interventions for elevated blood pressure and hypertension. *Nat Rev Cardiol.* 2021; 18: 785–802. DOI: <https://doi.org/10.1038/s41569-021-00559-8>
- Zhou B, Carrillo-Larco RM, Danaei G, Riley LM, Paciorek CJ, Stevens GA, Gregg EW, et al.** Worldwide trends in hypertension prevalence and progress in treatment and control from 1990 to 2019: A pooled analysis of 1201 population-representative studies with 104 million participants. *Lancet.* 2021; 398(10304): 957–980. DOI: [https://doi.org/10.1016/S0140-6736\(21\)01330-1](https://doi.org/10.1016/S0140-6736(21)01330-1)
- Mills KT, Bundy JD, Kelly TN, Reed JE, Kearney PM, Reynolds K, Chen J, He J.** Global disparities of hypertension prevalence and control: A systematic analysis of population-based studies from 90 countries. *Circulation.* 2016; 134(6): 441–450. DOI: <https://doi.org/10.1161/CIRCULATIONAHA.115.018912>
- Mills KT, Stefanescu A, He J.** The global epidemiology of hypertension. *Physiol Behav.* 2016; 176(1): 139–148.
- Chow CK, Teo KK, Rangarajan S, Islam S, Gupta R, Avezum A, Bahonar A, et al.** Prevalence, awareness, treatment, and control of hypertension in rural and urban communities in high-, middle-, and low-income countries. *J Am Med Assoc.* 2013; 310(9): 959–968. DOI: <https://doi.org/10.1001/jama.2013.184182>
- World Health Organization.** Indonesia: Noncommunicable diseases country profiles 2018 WHO. Geneva; 2018. <https://www.who.int/nmh/publications/ncd-profiles-2018/en/> Accessed: December 21, 2021.
- Kemendes RI.** Hasil Riset Kesehatan Dasar Tahun 2018. Vol. 53, Kementerian Kesehatan RI. Jakarta; 2018.
- Yusuf S, Rangarajan S, Teo K, Islam S, Li W, Liu L, Bo J, et al.** Cardiovascular risk and events in 17 low-, middle-, and high-income countries. *N Engl J Med.* 2014; 371(9): 818–827. DOI: <https://doi.org/10.1056/NEJMoa1311890>
- Ghembaza MA, Senoussaoui Y, Tani M, Meguenni K.** Impact of patient knowledge of hypertension complications on adherence to antihypertensive therapy. *Curr Hypertens Rev.* 2014; 10(1): 41–48. DOI: <https://doi.org/10.2174/157340211001141111160653>
- Turner BJ, Hollenbeak C, Weiner MG, Have TT, Roberts C.** Barriers to adherence and hypertension control in a racially diverse representative sample of elderly primary care patients. *Pharmacoepidemiol Drug Saf.* 2009; 18(5): 672–681. DOI: <https://doi.org/10.1002/pds.1766>
- Gheorghe A, Griffiths U, Murphy A, Legido-Quigley H, Lamptey P, Perel P.** The economic burden of cardiovascular disease and hypertension in low- and middle-income countries: A systematic review. *BMC Public Health.* 2018; 18(1): 1–11. DOI: <https://doi.org/10.1186/s12889-018-5806-x>
- Finkelstein EA, Chay J, Bajpai S.** The economic burden of self-reported and undiagnosed cardiovascular diseases and diabetes on Indonesian households. *PLoS One.* 2014; 9(6): e99572. DOI: <https://doi.org/10.1371/journal.pone.0099572>
- Kardas P, Lewek P, Matyjaszczyk M.** Determinants of patient adherence: A review of systematic reviews. *Front Pharmacol.* 2013 July; 4: 1–16. DOI: <https://doi.org/10.3389/fphar.2013.00091>
- Maimaris W, Paty J, Perel P, Legido-Quigley H, Balabanova D, Nieuwlaat R, Mckee M.** The influence of health systems on hypertension awareness, treatment, and control: A systematic literature review. *PLoS Med.* 2013; 10(7): e1001490. DOI: <https://doi.org/10.1371/journal.pmed.1001490>
- Sujarwoto S, Maharani A.** Participation in community-based health care interventions (CBHIs) and its association with hypertension awareness, control and treatment in Indonesia. *PLoS One.* 2020; 15(12): 1–18. DOI: <https://doi.org/10.1371/journal.pone.0244333>
- Felix C, Baldeon ME, Zertuche F, Fornasini M, Paucar MJ, Ponce L, Rangarajan S, Yusuf S, Lopez-Jaramillo P.** Low levels of awareness, treatment, and control of hypertension in Andean communities of Ecuador. *J Clin Hypertens.* 2020; 22(9): 1530–1537. DOI: <https://doi.org/10.1111/jch.13982>

17. **Peltzer K, Pengpid S.** The prevalence and social determinants of hypertension among adults in Indonesia: A cross-sectional population-based national survey. *Int J Hypertens*; 2018. DOI: <https://doi.org/10.1155/2018/5610725>
18. **Cifkova R, Fodor G, Wohlfahrt P.** Changes in hypertension prevalence, awareness, treatment, and control in high-, middle-, and low-income countries: An update. *Curr Hypertens Rep.* 2016; 18(8): 1–6. DOI: <https://doi.org/10.1007/s11906-016-0669-y>
19. **Strauss J, Witoelar F, Sikoki B.** The fifth wave of the Indonesia Family Life Survey: Overview and field report: Volume 1. *Fifth Wave Indones Fam Life Surv Overv F Rep.* 2016 March; 1: 1–94. DOI: <https://doi.org/10.7249/WR1143.1>
20. **Hussain MA, Al Mamun A, Reid C, Huxley RR.** Prevalence, awareness, treatment and control of hypertension in Indonesian adults aged ≥ 40 years: Findings from the Indonesia Family Life Survey (IFLS). *PLoS One.* 2016; 11(8): 1–16. DOI: <https://doi.org/10.1371/journal.pone.0160922>
21. **James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, Lackland DT,** et al. 2014 evidence-based guideline for the management of high blood pressure in adults: Report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *J Am Med Assoc.* 2014; 311(5): 507–520. DOI: <https://doi.org/10.1001/jama.2013.284427>
22. **McAlister FA, Wilkins K, Joffres M, Leenen FHH, Fodor G, Gee M, Tremblay MS, Walker R, Johansen H, Campbell N.** Changes in the rates of awareness, treatment and control of hypertension in Canada over the past two decades. *CMAJ.* 2011; 183(9): 1007–1013. DOI: <https://doi.org/10.1503/cmaj.101767>
23. **Damasceno A, Azevedo A, Silva-Matos C, Prista A, Diogo D, Lunet N.** Hypertension prevalence, awareness, treatment, and control in mozambique: Urban/rural gap during epidemiological transition. *Hypertension.* 2009; 54(1): 77–83. DOI: <https://doi.org/10.1161/HYPERTENSIONAHA.109.132423>
24. **Vrijens B, De Geest S, Hughes DA, Przemyslaw K, Démonceau J, Ruppert T, Dobbels F,** et al. A new taxonomy for describing and defining adherence to medications. *Br J Clin Pharmacol.* 2012; 73(5): 691–705. DOI: <https://doi.org/10.1111/j.1365-2125.2012.04167.x>
25. **Vilagut G, Forero CG, Barbaglia G, Alonso J.** Screening for depression in the general population with the center for epidemiologic studies depression (ces-d): A systematic review with meta-analysis. *PLoS One.* 2016; 11(5): 1–17. DOI: <https://doi.org/10.1371/journal.pone.0155431>
26. **Björgvinsson T, Kertz SJ, Bigda-Peyton JS, McCoy KL, Aderka IM.** Psychometric properties of the CES-D-10 in a psychiatric sample. *Assessment.* 2013; 20(4): 429–436. DOI: <https://doi.org/10.1177/1073191113481998>
27. **Yu L, Buysse DJ, Germain A, Moul DE, Stover A, Dodds NE, Johnston KL, Piconis PA.** Development of short forms from the PROMIS Sleep Disturbance and Sleep-Related Impairment item banks. *Behav Sleep Med.* 2011; 10(1): 6–24. DOI: <https://doi.org/10.1080/15402002.2012.636266>
28. **Peltzer K, Pengpid S.** Prevalence, social and health correlates of insomnia among persons 15 years and older in Indonesia. *Psychol Heal Med.* 2019; 24(6): 757–768. DOI: <https://doi.org/10.1080/13548506.2019.1566621>
29. **IFLS-5.** Buku Iii B. In: *Survei Aspek Kehidupan Rumah Tangga Indonesia 2014.* Jakarta: RAND; 2014.
30. **Dodge Y.** Chi-square goodness of fit test. In: *The concise encyclopedia of statistics.* New York: Springer. 2008; pp. 72–76. DOI: https://doi.org/10.1007/978-0-387-32833-1_55
31. **Berkman ND, Sheridan SL, Donahue KE, Halpern DJ, Crotty K.** Low health literacy and health outcomes: An updated systematic review. *Ann Intern Med.* 2011; 155(2): 97–107. DOI: <https://doi.org/10.7326/0003-4819-155-2-201107190-00005>
32. **Mekonnen GB, Gelayee DA.** Low medication knowledge and adherence to oral chronic medications among patients attending community pharmacies: A cross-sectional study in a low-income country. *Biomed Res Int.* 2020. DOI: <https://doi.org/10.1155/2020/4392058>
33. **Gordon K, Smith F, Dhillon S.** Effective chronic disease management: Patients' perspectives on medication-related problems. *Patient Educ Couns.* 2007; 65(3): 407–415. DOI: <https://doi.org/10.1016/j.pec.2006.09.012>
34. **Cutler RL, Fernandez-Llimos F, Frommer M, Benrimoj C, Garcia-Cardenas V.** Economic impact of medication non-adherence by disease groups: A systematic review. *BMJ Open.* 2018; 8(1): e016982. DOI: <https://doi.org/10.1136/bmjopen-2017-016982>
35. **Ip Q, Malone DC, Chong J, Harris RB, Labiner DM.** Economic impact of epilepsy and the cost of nonadherence to antiepileptic drugs in older Medicare beneficiaries. *Epilepsy Behav.* 2018; 80: 208–214. DOI: <https://doi.org/10.1016/j.yebeh.2018.01.009>
36. **Desai R, Nayak R.** Effects of medication nonadherence and comorbidity on health resource utilization in schizophrenia. *J Manag Care Spec Pharm.* 2019; 25(1): 37–44A. DOI: <https://doi.org/10.18553/jmcp.2019.25.1.037>
37. **Alfian SD, Sukandar H, Lestari K, Abdulah R.** Medication adherence contributes to an improved quality of life in type 2 diabetes mellitus patients: A cross-sectional study. *Diabetes Ther Res Treat Educ Diabetes Relat Disord.* 2016; 7(4): 755–764. DOI: <https://doi.org/10.1007/s13300-016-0203-x>
38. **Lee JH, Kim K-IL, Cho MC.** Current status and therapeutic considerations of hypertension in the elderly. *Korean J Intern Med.* 2019; 34(4): 687–695. DOI: <https://doi.org/10.3904/kjim.2019.196>

39. **Kretchy I, Owusu-Daaku F, Danquah S.** Spiritual and religious beliefs: Do they matter in the medication adherence behaviour of hypertensive patients? *Biopsychosoc Med.* 2013; 7(1): 1. DOI: <https://doi.org/10.1186/1751-0759-7-15>
40. **Shahin W, Kennedy GA, Stupans I.** The impact of personal and cultural beliefs on medication adherence of patients with chronic illnesses: A systematic review. *Patient Prefer Adherence.* 2019; 13: 1019–1035. DOI: <https://doi.org/10.2147/PPA.S212046>
41. **Albuquerque UP, Ferreira Júnior WS, Sousa DCP, Reinaldo RCPS, do Nascimento ALB, Gonçalves PHS.** Religiosity/spirituality matters on plant-based local medical system. *J Relig Health.* 2018; 57(5): 1948–1960. DOI: <https://doi.org/10.1007/s10943-018-0634-y>
42. **Frazão-Moreira A.** The symbolic efficacy of medicinal plants: Practices, knowledge, and religious beliefs amongst the Nalu healers of Guinea-Bissau. *J Ethnobiol Ethnomed.* 2016; 12(1): 1–15. DOI: <https://doi.org/10.1186/s13002-016-0095-x>
43. **Asfaw Erku D, Basazn Mekuria A.** Prevalence and correlates of complementary and alternative medicine use among hypertensive patients in Gondar Town, Ethiopia. *Evidence-Based Complement Altern Med.* 2016. DOI: <https://doi.org/10.1155/2016/6987636>
44. **Alfian SD, Sukandar H, Arisanti N, Abdulah R.** Complementary and alternative medicine use decreases adherence to prescribed medication in diabetes patients. *J Acupunct Meridian Stud.* 2018; 11(5): 337. DOI: <https://doi.org/10.1016/j.jams.2018.10.002>
45. **Li H, Yan X, Deng X, Yang L, Zhao S, Zou J, Luo Y, Cao S.** A cross-sectional study of the prevalence, awareness, treatment and control of hypertension in Shenzhen, China. *BMJ Open.* 2017; 7(6): 1–7. DOI: <https://doi.org/10.1136/bmjopen-2016-015206>
46. **Turana Y, Teng kawan J, Soenarta AA.** Asian management of hypertension: Current status, home blood pressure, and specific concerns in Indonesia. *J Clin Hypertens.* 2020; 22(3): 483–485. DOI: <https://doi.org/10.1111/jch.13681>
47. **Kiselev AR, Posnenkova OM, Belova OA, Romanchuk SV, Popova YV, Prokhorov MD, Gridnev VI.** Impact of clinical factors on the achievement of target blood pressure in hypertensive patients from Ivanovo Region of Russia: Data of 2015. *High Blood Press Cardiovasc Prev.* 2017; 24(4): 425–435. DOI: <https://doi.org/10.1007/s40292-017-0227-y>
48. **Carey RM, Muntner P, Bosworth HB, Whelton PK.** Prevention and control of hypertension: JACC Health Promotion Series. *J Am Coll Cardiol.* 2018; 72(11): 1278–1293. DOI: <https://doi.org/10.1016/j.jacc.2018.07.008>
49. **Odedosu T, Schoenthaler A, Vieira DL, Agyemang C, Ogedegbe G.** Overcoming barriers to hypertension control in African Americans. *Cleve Clin J Med.* 2012; 79(1): 46–56. DOI: <https://doi.org/10.3949/ccjm.79a.11068>
50. **Jneid S, Jabbour H, Hajj A, Sarkis A, Licha H, Hallit S, Khabbaz LR.** Quality of life and its association with treatment satisfaction, adherence to medication, and trust in physician among patients with hypertension: A cross-sectional designed study. *J Cardiovasc Pharmacol Ther.* 2018; 23(6): 532–542. DOI: <https://doi.org/10.1177/1074248418784292>
51. **Świątoniowska-Lonc N, Polański J, Tański W, Jankowska-Polańska B.** Impact of satisfaction with physician-patient communication on self-care and adherence in patients with hypertension: Cross-sectional study. *BMC Health Serv Res.* 2020; 20(1): 1–9. DOI: <https://doi.org/10.1186/s12913-020-05912-0>
52. **Botha E, Gwin T, Purpora C.** The effectiveness of mindfulness based programs in reducing stress experienced by nurses in adult hospital settings: a systematic review of quantitative evidence protocol. *JBI database Syst Rev Implement reports.* 2015; 13(10): 21–29. DOI: <https://doi.org/10.11124/jbisrir-2015-2380>
53. **Zhang Y, Chen Y, Ma L.** Depression and cardiovascular disease in elderly: Current understanding. *J Clin Neurosci.* 2018; 47: 1–5. DOI: <https://doi.org/10.1016/j.jocn.2017.09.022>
54. **Meng L, Chen D, Yang Y, Zheng Y, Hui R.** Depression increases the risk of hypertension incidence: A meta-analysis of prospective cohort studies. *J Hypertens.* 2012; 30(5): 842–851. DOI: <https://doi.org/10.1097/HJH.0b013e32835080b7>
55. **Hare DL, Toukhsati SR, Johansson P, Jaarsma T.** Depression and cardiovascular disease: A clinical review. *Eur Heart J.* 2014; 35(21): 1365–1372. DOI: <https://doi.org/10.1093/eurheartj/eh462>
56. **Read JR, Sharpe L, Modini M, Dear BF.** Multimorbidity and depression: A systematic review and meta-analysis. *J Affect Disord.* 2017; 221: 36–46. DOI: <https://doi.org/10.1016/j.jad.2017.06.009>
57. **Lian Y, Zhang J, Jia CX.** Sleep duration change and cognitive function: A national cohort study of Chinese people older than 45 years. *J Nerv Ment Dis.* 2020; 208(6): 498–504. DOI: <https://doi.org/10.1097/NMD.0000000000001159>
58. **Hua J, Sun H, Shen Y.** Improvement in sleep duration was associated with higher cognitive function: a new association. *Aging (Albany NY).* 2020; 12(20): 20623–20635. DOI: <https://doi.org/10.18632/aging.103948>
59. **Unger T, Borghi C, Charchar F, Khan NA, Poulter NR, Prabhakaran D, Ramirez A, et al.** 2020 International Society of Hypertension Global Hypertension practice guidelines. *Hypertension.* 2020; 75(6): 1334–1357. DOI: <https://doi.org/10.1161/HYPERTENSIONAHA.120.15026>

60. **Odili AN, Chori BS, Danladi B, Nwakile PC, Okoye IC, Abdullahi U, Nwegbu MN, et al.** Prevalence, awareness, treatment and control of hypertension in Nigeria: Data from a nationwide survey 2017. *Glob Heart*. 2020; 15(1): 1–13. DOI: <https://doi.org/10.5334/gh.848>
61. **Mirzaei M, Mirzaei M, Bagheri B, Dehghani A.** Awareness, treatment, and control of hypertension and related factors in adult Iranian population. *BMC Public Health*. 2020; 20(1): 1–10. DOI: <https://doi.org/10.1186/s12889-020-08831-1>
62. **Olives C, Myerson R, Mokdad AH, Murray CJL, Lim SS.** Prevalence, awareness, treatment, and control of hypertension in United States Counties, 2001–2009. *PLoS One*. 2013; 8(4): e60308. DOI: <https://doi.org/10.1371/journal.pone.0060308>
63. **Yip W, Wong TY, Jonas JB, Zheng Y, Lamoureux EL, Nangia V, Sabanayagam C.** Prevalence, awareness, and control of hypertension among Asian Indians living in urban Singapore and rural India. *J Hypertens*. 2013; 31(8): 1539–1546. DOI: <https://doi.org/10.1097/HJH.0b013e328361d52b>
64. **Sohn K.** Sick but unaware: Hypertension in Indonesia. *Soc Biodemography Soc Biol*. 2015; 61(3): 298–318. DOI: <https://doi.org/10.1080/19485565.2015.1091719>
65. **Guimond AJ, Kubzansky LD, Boehm JK, Kivimaki M, Trudel-Fitzgerald C.** Does life satisfaction reduce risk of incident hypertension and stroke? Evidence from the Whitehall II cohort. *J Psychosom Res*. 2021; 144: 1–9. November 2020. DOI: <https://doi.org/10.1016/j.jpsychores.2021.110414>
66. **Christiani Y, Byles JE, Tavener M, Dugdale P.** Health insurance coverage among women in Indonesia's major cities: A multilevel analysis. *Health Care Women Int*. 2017; 38(3): 267–282. DOI: <https://doi.org/10.1080/07399332.2016.1253697>
67. **Gebru T, Lentiro K.** The impact of community-based health insurance on health-related quality of life and associated factors in Ethiopia: A comparative cross-sectional study. *Health Qual Life Outcomes*. 2018; 16(1): 1–6. DOI: <https://doi.org/10.1186/s12955-018-0946-3>
68. **Romain M, Sprung CL.** Approaches to patients and families with strong religious beliefs regarding end-of-life care. *Curr Opin Crit Care*. 2014; 20(6): 668–672. DOI: <https://doi.org/10.1097/MCC.000000000000148>
69. **Aulia G, Halimah E, Lestari K.** Pharmaceutical counseling has a positive impact on quality of life of hypertension patients. *Pharmacol Clin Pharm Res*. 2018; 3(1): 5–9. DOI: <https://doi.org/10.15416/pcpr.v3i1.16449>
70. **Graham B.** Defining and measuring patient satisfaction. *J Hand Surg Am*. 2016; 41(9): 929–931. DOI: <https://doi.org/10.1016/j.jhssa.2016.07.109>
71. **Alfian SD, Annisa N, Fajriansyah F, Perwitasari DA, Abdulah R, Hak E, Denig P.** Modifiable factors associated with non-adherence to antihypertensive or antihyperlipidemic drugs are dissimilar: A multicenter study among patients with diabetes in Indonesia. *J Gen Intern Med*. 2020; 35(10): 2897–2906. DOI: <https://doi.org/10.1007/s11606-020-05809-y>
72. **Fenech G, Vallée A, Cherfan M, Kab S, Goldberg M, Zins M, Blacher J.** Poor awareness of hypertension in France: The CONSTANCES population-based study. *Am J Hypertens*. 2020; 33(6): 543–551. DOI: <https://doi.org/10.1093/ajh/hpaa018>
73. **Villalva CM, Alvarez-Muiño XLL, Mondelo TG, Fachado AA, Fernández JC.** Adherence to treatment in hypertension. *Adv Exp Med Biol*. 2017; 956: 129–147. DOI: https://doi.org/10.1007/5584_2016_77

TO CITE THIS ARTICLE:

Khoiry QA, Alfian SD, Abdulah R. Modifiable and Non-modifiable Factors Associated with Low Awareness of Hypertension Treatment in Indonesia: A Cross-Sectional Population-Based National Survey. *Global Heart*. 2022; 17(1): 56. DOI: <https://doi.org/10.5334/gh.1143>

Submitted: 28 April 2022

Accepted: 26 July 2022

Published: 16 August 2022

COPYRIGHT:

© 2022 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.

Global Heart is a peer-reviewed open access journal published by Ubiquity Press.