

ORIGINAL RESEARCH

Ideal Cardiovascular Health in a Nationally Representative Population-Based Sample of Adults in Malawi

Supa Pengpid^{1,2} and Karl Peltzer^{3,4}¹ ASEAN Institute for Health Development, Mahidol University, Salaya, Phutthamonthon, Nakhon Pathom, TH² Department of Research Administration and Development, University of Limpopo, Turfloop, ZA³ Department of Psychology, University of the Free State, Bloemfontein, ZA⁴ Department of Psychology, College of Medical and Health Science, Asia University, Taichung, TW

Corresponding author: Karl Peltzer (kfpeltzer@gmail.com)

Background: Ideal cardiovascular health (ICH) measures four ideal health behaviours (non-smoking, body mass index <25 kg/m², healthy diet, and physical activity) and three health factors (total cholesterol <200 mg/dL, blood pressure <120/<80 mmHg, and fasting blood glucose <100 mg/dL).

Objective: This study aimed to determine the prevalence, distribution, and correlates of ICH among adults in Malawi.

Methods: National cross-sectional survey data of 3,441 persons aged 18–69 years with complete ICH measurements in Malawi in 2017 were analysed.

Results: Almost one in ten (7.4% of respondents had 0–2 ICH metrics), 21.2% 3–4 ICH metrics, and 71.5% 5–7 ICH metrics). Only 3.3% had all seven ICH metrics, 15.3% had intermediate ICH (≥1 metric in the intermediate category and none in the poor category), and 81.5% poor ICH (≥1 metric in poor category). In adjusted logistic regression analysis, older age (50–69 years) (Adjusted Odds Ratio-AOR: 0.25, 95% Confidence Interval-CI: 0.17–0.36) and urban residence (AOR: 0.56, 95% CI: 0.40–0.78) were negatively associated with meeting 5–7 ICH metrics. In addition, in unadjusted analysis, higher education was positively associated with meeting 5–7 ICH metrics.

Conclusion: The proportion of meeting 5–7 ICH metrics was high in Malawian adults. Both high-risk and population-wide intervention programmes targeting older adults and urban residents should be implemented in aiding to improve cardiovascular health in Malawi.

Highlights

- The proportion of meeting 5–7 ideal cardiovascular health (ICH) metrics was 71.5% in Malawian adults (18–69 years, median age 32 years).
- In a sub-group analysis (45–69 years, median age 55 years), 55.9% had 5–7 ideal ICH metrics.
- ICH was lower with increasing age and among urban residents.

Keywords: cardiovascular health; adults; Malawi

Background

Globally, it is estimated that 31% of all death are attributed to cardiovascular diseases (CVDs), such as heart attacks and stroke, in 2016 [1]. Ischaemic heart disease and stroke were the major causes of disability-adjusted life years (DALYs) in persons 50 years and older in 2019 [2]. More than three-quarters of deaths from CVDs occur in low- and middle-income countries [1].

To reduce CVDs, the American Heart Association (AHA) developed the concept of ideal cardiovascular health (ICH), including seven ideal health behaviours and factors [3, 4]. An increasing number of ICH metrics have been found protective against 'the all-cause and CVD-related mortality risk, incident cardiovascular events, lower prevalence and incidence of non-CVD outcomes such as cancer, depression, and cognitive

impairment' [5]. To our knowledge, there are no national data on ICH in Malawi, a low-income country in Southern Africa. CVDs contribute to 10% of mortality in 2016 in Malawi [6]. In Malawi, a high burden of CVD risk factors has been found, including hypertension and diabetes, smoking, heavy alcohol use, physical inactivity, high salt and sugar intake, low fruit and vegetable intake, and obesity [7, 8]. CVD is on the rise in Africa, mainly attributable to an increase in hypertension, smoking, and obesity [9].

Globally, mainly in high-income countries, 32.2% (having 0–2 ideal metrics) had overall poor and 19.6% of participants had 5–7 ICH metrics [10]. Fewer studies have been conducted on ICH in Africa. In rural South Africa (46% HIV positive, 38.7 years mean age), 7% had poor (0–2 metrics), 40% intermediate (3–4 metrics) and 53% had 5–7 ICH metrics [11], in Uganda (N = 857, ≥ 16 years, mean age 41.5 years in women and 36.8 years in men), 3.2% had 7 ICH metrics, 50.0% had 5–7 ICH metrics [12], and in urban (n = 596) and rural (n = 326) Ghana (25–70 years, mean age 44 and 45 years in rural and urban women, respectively), urban women 74.7% ≥ 1 metric in poor category, 25.0% in ≥ 1 metric in the intermediate category and none in the poor category, 0.3% all 7 ICH metrics [13]. Globally, smoking had the highest prevalence of ICH status (69.1%), followed by fasting blood glucose (FBG) (67.7%), total cholesterol (TC) (51.7%), physical activity (PA) (40.6%), body mass index (BMI) (40.3%), blood pressure (BP) (34.6%), and dietary pattern (12.1%) [10].

Sociodemographic factors associated with ICH may include female sex [63 high income countries-HIC and 28 low- and middle-income countries-LMIC, 10, Brazil, 14, Nepal, 15], younger age [63 HIC and 28 LMIC, 10, Uganda, 12, Brazil, 14, Nepal, 15, China, 16], ethnicity [Brazil, 14], higher education [Uganda, 12, Brazil, 14], higher income [Jamaica, 17, China, 18], lower income [Uganda, 12], rural residence [Ghana, 13, Peru, 19], and geographic region [Brazil, 20]. This study aimed to determine the prevalence, distribution, and correlates of ideal ICH among adults in Malawi in 2017.

Method

Sample and procedures

The study design was a national community based cross-sectional survey in Malawi in 2017, using WHO STEPwise approach for assessing risk factors for chronic non-communicable diseases (NCDs), and includes questionnaire data on demographic and behavioural information (Step 1), physical measurements (Step 2) and biochemistry tests (Step 3) [21]. Based on a multistage cluster sample design, the '2017 Malawi STEPS Survey' produced nationally representative population-based data for persons 18–69 years in Malawi [21]. Inclusion criteria was one randomly selected adult household member aged 18–69 per household who was able to provide informed consent. More information on the study methods and data can be publicly accessed [21]. The inclusion criteria for the present study were individuals with no data missing for smoking status, BMI, PA, diet, total TC, FBG, and BP measurements. From 4,187 adults, a total of 3,441 participants with full required information were included. Ethics approval was obtained from the 'Malawi National Ethics Committee' and participants provided written informed consent [21].

Measures

Poor, intermediate and ICH levels for smoking, BMI, PA, diet, TC, BP, and FBG were determined, based on modified AHA definitions [3, 4], exact AHA definitions are provided in brackets.

Cardiovascular health behaviour

Smoking status: Smoking is defined as poor if current smoker, intermediate [former ≤ 12 months] if a past smoker, and ideal if self-report of [never or quit > 12 months] never having smoked.

Body Mass Index (BMI) (kg/m²): BMI is defined poor if ≥ 30 kg/m², intermediate as 25.0–29.9 kg/m², and ideal BMI is < 25 kg/m². Anthropometric measurements were taken by trained healthcare staff using a portable electronic weighing scale and measuring inflexible bars [21].

Healthy diet: [poor: 0–1 components, intermediate: 2–3, and ideal: 4–5 components (1: ≥ 4.5 cups/day fruits and vegetables, 2: ≥ 3.5 ounce servings/week of fish, 3: < 1500 milligrams/day sodium, 4: < 450 calories/week sweets/sugar, and 5: ≥ 3 1-ounce servings/day whole grains)]. Poor healthy diet is defined in this study as < 2 servings of fruit and vegetables (FV)/day, intermediate as 2– < 4.5 FV/day, and an ideal diet as ≥ 4.5 FV servings/day.

Physical activity (PA): 'Poor = None, Intermediate = 1–149 min/wk moderate intensity or 1–74 min/wk vigorous intensity or 1–149 min/wk moderate+vigorous, ideal = ≥ 150 min/wk moderate intensity or ≥ 75 min/wk vigorous intensity or ≥ 150 min/wk moderate+vigorous.' PA was assessed with the Global Physical Activity Questionnaire [22].

Cardiovascular health factors

Poor total cholesterol (TC) is classified as 'TC ≥ 6.3 mmol/L (≥ 240 mg/dL), intermediate is TC 5.2–6.2 mmol/L (200–239 mg/dL) or treated to TC < 5.2 mmol/L (< 200 mg/dL) and ideal TC is < 200 mg/dL and without any cholesterol-lowering medication.'

Fasting blood glucose (FBG): poor FBG is defined as 'glucose ≥ 7.0 mmol/L (≥ 126 mg/dL), intermediate is glucose 5.6–5.9 mmol/L (100–125 mg/dL) or treated to < 100 mg/dL, and ideal is < 5.6 mmol/L < 100 mg/dL and without any glucose-lowering medication.'

For TC and FBG finger blood samples for biochemistry tests were taken, provided instructions of starving overnight were followed, and TV and FBG were measured using Cardiochek [Report].

Blood pressure (BP): poor is defined as 'BP $\geq 140/\geq 90$ mmHg, intermediate is systolic BP 120–139 mmHg or diastolic BP 80–89 mmHg or treated to BP $< 120/< 80$ mmHg, and ideal BP is defined as BP $< 120/< 80$ mmHg and without any antihypertensive medication.' Of the three BP measurements (taken 3–5 minutes apart) using digital BP machines (Omron M4-1), the last two readings were averaged [21].

The seven ICH items were dichotomised (1 = ideal, 0 = not ideal), and grouped into 0–2, 3–4, and 5–7 ICH metric; 5–7 ICH metric is in the absence of any previous CVD. In addition, three ICH groups were created as follows: 'ICH is all seven health metrics at ideal levels in the absence of any previous CVD, intermediate ICH is at least one health metric at the intermediate level, but no poor ICH metrics, and poor ICH is at least one of seven ICH metrics at poor level' [3, 4, 23]. Ideal health behaviour was defined as the simultaneous presence of 4 ideal health behaviours (adequate PA, nonsmoker, normal BMI, and healthy diet) and ideal health factors as the simultaneous presence of four ideal health factors (non-smokers, normal BP, normal FBG, and normal TC) [3, 4, 23].

History of CVDs was assessed with the question, 'Have you ever had a heart attack or chest pain from heart disease (angina) or a stroke (cerebrovascular accident or incident)?' (Yes, No) [21].

Sociodemographic covariates included age, sex, highest level of formal education, work, and residence status [21].

Data analysis

Considering the clustered study design, all statistical analyses were conducted with 'STATA software version 14.0 (Stata Corporation, College Station, TX, USA).' Only participants with complete ICH assessments were included in the analysis. Chi-square tests were used for estimating differences in proportions and Student's t-test for differences in means. ICH metrics are described across ideal, intermediate, and poor ICH. Unadjusted and adjusted logistic regressions were used to calculate sociodemographic predictors (age group, sex, educational level, employment status and residence status) of meeting 5–7 ICH metrics, overall and for men and women separately. P-values of below 0.05 were accepted as significant and missing values were excluded from the analysis.

Results

Sample characteristics

The sample included 3,441 adults (18–69 years), with a median age of 32 years (interquartile range 24–42), 35.9% were male. More than half of the participants (59.0%) Standard 5 or higher education, 42.9% were employed or students, and 90.0% lived in rural areas. The mean BMI of the respondents was 22.6, the mean systolic BP was 121.1 mmHg, and the prevalence of self-reported CVD was 6.8%. Compared to men, women had lower education, lower employed or student status, lower systolic BP, more likely living in urban areas, more likely having a CVD, more likely taking antihypertensive and lipid-lowering drugs, having a higher BMI, and higher total TC (see **Table 1**).

Distribution of cardiovascular health metrics

The distribution of the three levels of all seven ICH metrics in the overall adult population and by sex is shown in **Table 2**. Approximately, 84.0% of Malawian adults reported that they never smoked (70.9% in men and 97.4% in women). More than four in five participants (81.8%) had ideal BMI (90.7% in men and 72.8% in women), and 97.8% had ideal physical activity (99.0% in men and 96.7% in women). A low proportion of healthy diet (≥ 4.5 servings of fruit and vegetables/day) of 11.3% was reported (9.7% among men and 13.0% among women). Most Malawian adults had ideal total cholesterol (95.0%) and fasting glucose levels (92.7%), while only 46.9% had ideal blood pressure. A significant higher proportion of women had ideal smoking and ideal blood pressure than men, while men had significantly higher ideal BMI, PA and TC than women. Almost one in ten (7.4%) of respondents had poor ICH (0–2 ideal metrics),

Table 1: Sample characteristics of participants aged 18-69, Malawi, 2017.

Variable	Total (N = 3441)	Men (n = 1237)	Women (n = 2204)
	%	%	%
All		35.9	64.1
Age (years)			
18–29	44.9	47.5	42.2
30–49	39.8	38.4	41.3
50–69	15.3	14.1	16.6
Education			
Secondary or more	23.5	26.1	20.9*
Standard 5–8	35.5	40.4	30.6
Standard 1–4	31.1	28.1	34.2
None	9.9	5.5	14.3
Employment status			
Nonpaid or unemployed	57.1	71.4	43.2*
Employed or student	42.9	28.6	56.8
Residence			
Rural	90.0	92.0	87.9*
Urban	10.0	8.0	12.1
Self-reported cardiovascular disease	6.8	4.6	9.1*
Use of anti-hypertensive drug	2.4	0.8	4.0*
Use of hypoglycemic drug	0.3	0.4	0.2
Use of lipid-lowering drug	0.1	0.0	0.1*
	M (SD)	M (SD)	M (SD)
Mean systolic blood pressure, mmHg	121.1 (15.9)	122.4 (15.4)*	119.7 (16.2)
Mean body mass index, kg/m ²	22.6 (4.0)	21.7 (3.2)	23.6 (4.4)*
Mean total cholesterol, mmol/L	3.5 (0.9)	3.3 (0.8)	3.7 (0.9)*
Mean fasting plasma glucose, mmol/L	4.7 (1.0)	4.6 (1.4)	4.8 (1.1)*

* $p < 0.05$, men versus women.

21.2% intermediate ICH (3–4 ideal metrics), and 71.5% ideal ICH (5–7 ideal metrics). In a sub-group analysis, persons aged 45–69 years, 55.9% had 5–7 ICH metrics. Only 3.3% had all seven ICH metrics, 15.3% intermediate ICH (≥ 1 metric in the intermediate category and none in the poor category), and 81.5% had poor ICH (≥ 1 metric in poor category). Men had better ICH metrics than women in the first ICH measure and women had better ICH than men in the second measure. Compared to the 18- to 44-year-olds, 45- to 69-year-olds had poorer overall ICH metrics as well as six individual ICH, except for fruit and vegetable intake (see **Table 2**).

Proportion of ideal cardiovascular health metrics

In all, 0.0% had zero, 0.1% one, 0.6% two, 4.5% three, 18.6% four, 40.3% five, 32.3% six and 3.6% all seven ICH metrics (see **Table 3**). A total of 36.1% participants were ideal on all four health factors, but only 7.3% were ideal on all four health behaviours, the proportion of all four health factors was significantly higher among women (44.2%) than men (28.1%) ($p < 0.001$) as well as for all four health behaviours (9.0% among women and 5.6% among men) ($p < 0.001$).

Table 2: Ideal cardiovascular health (ICH) metrics distribution (prevalence, %).

Health metrics		Total sample (N = 3441)	Men (n = 1237)	Women (n = 2204)	Chi-square p-value, Men vs Women	Age: 18–44 years	Age: 45–69 years	Chi-square p-value, 18–44 vs 45–69
Smoking	Poor	11.4	20.9	1.7	<0.001	9.9	16.4	<0.001
	Intermediate	4.5	8.1	0.9		2.9	9.4	
	Ideal	84.0	70.9	97.4		87.2	74.2	
Body mass index	Poor	5.0	1.0	9.1	<0.001	4.4	6.7	0.008
	Intermediate	13.2	8.3	18.1		12.3	16.5	
	Ideal	81.8	90.7	72.8		83.3	76.7	
Diet: Fruit and vegetable intake	Poor	71.3	74.8	67.8	0.066	72.2	72.3	0.833
	Intermediate	17.4	15.5	19.3		16.8	17.4	
	Ideal	11.3	9.7	13.0		11.1	10.3	
Physical activity	Poor	1.1	0.5	1.6	0.007	0.7	2.3	<0.001
	Intermediate	1.1	0.5	1.7		0.9	1.7	
	Ideal	97.8	99.0	96.7		98.4	96.1	
Total cholesterol	Poor	0.7	0.2	1.2	<0.001	0.5	1.9	<0.001
	Intermediate	4.2	2.5	6.0		3.2	9.0	
	Ideal	95.0	97.3	92.8		96.3	89.1	
Blood pressure	Poor	15.6	14.9	16.3	0.004	11.0	30.9	<0.001
	Intermediate	37.5	42.1	32.9		38.8	32.1	
	Ideal	46.9	43.0	50.8		50.2	37.1	
Fasting plasma glucose	Poor	1.5	1.4	1.6	0.307	1.3	1.7	<0.001
	Intermediate	5.9	4.7	7.0		4.5	9.3	
	Ideal	92.7	93.9	91.4		94.2	89.0	
0–2 ICH metrics		7.4	4.9	9.9	<0.001	5.9	12.9	<0.001
3–4 ICH metrics		21.2	22.3	20.0		18.4	31.2	
5–7 ICH metrics)		71.5	72.8	70.2		75.7	55.9	
Ideal ICH ^a		3.3	1.6	4.9	<0.001	3.9	1.0	<0.001
Intermediate ICH ^b		15.3	13.5	17.1		16.3	11.4	
Poor ICH ^c		81.5	84.8	78.1		79.8	87.6	

^a All seven ICH metrics at ideal levels in the absence of cardiovascular disease (CVD), ^b At least one of seven ICH metrics at intermediate levels, no poor ICH metrics in participants without CVD history or if all seven ICH metrics are ideal among persons with a CVD history, ^c At least one of seven ICH metrics at a poor level in participants without a CVD history or at least one metric is intermediate or poor among persons with a CVD history' [10, 25].

Associations with meeting 5–7 ICH metrics

In adjusted logistic regression analysis, older age (50–69 years) (Adjusted Odds Ratio-AOR: 0.25, 95% Confidence Interval-CI: 0.17–0.36) and urban residence (AOR: 0.56, 95% CI: 0.40–0.78) were negatively associated with meeting 5–7 ideal CVH metrics. In addition, in unadjusted analysis, higher education was positively associated with meeting 5–7 ICH metrics (see **Table 4**).

Discussion

To our knowledge, this is the first national study on the prevalence and distribution of ICH metrics in Malawi. In this nationally representative sample in Malawi, the prevalence of poor ICH (0–2 ideal metrics) (7.4%), and 5–7 ICH metrics (71.5%) in the total adult population (18–69 years, median age 32 years) and poor

Table 3: Distribution of ideal cardiovascular health (ICH) metrics in percent among participants.

Variable	Sample	Proportion of ICH metrics							
	N	0	1	2	3	4	5	6	7
All	3441	0.0	0.1	0.6	4.5	18.6	40.3	32.3	3.6
Age in years									
18–29	1037	0.0	0.0	0.0	2.2	12.9	40.5	40.0	4.4
30–49	1588	0.0	0.1	0.7	4.9	21.0	41.1	28.6	3.5
50–69	816	0.0	0.6	1.9	10.3	28.9	37.3	19.8	1.3
45–69	1108	0.0	0.4	1.9	8.7	26.5	38.4	23.0	1.0
Sex									
Female	2204	0.0	0.1	0.8	4.7	18.3	35.7	35.0	5.3
Male	1237	0.0	0.1	0.4	4.3	18.8	44.8	29.7	1.9
Education									
Secondary or more	861	0.0	0.2	0.7	3.8	17.3	45.1	31.0	1.9
Standard 5–8	1023	0.0	0.0	0.5	3.6	14.7	39.5	37.4	4.3
Standard 1–4	1073	0.0	0.1	0.4	4.7	22.4	38.5	29.9	4.1
None	481	0.0	0.6	1.2	9.2	23.6	37.2	24.8	3.4
Employment status									
Nonpaid or unemployed	1963	0.0	0.2	0.6	4.5	18.1	38.5	33.5	4.6
Employed or student	2204	0.0	0.0	0.6	4.6	19.2	42.6	30.8	2.3
Residence									
Rural	2760	0.0	0.1	0.4	4.3	17.9	40.2	33.5	3.7
Urban	681	0.0	0.3	2.5	6.5	24.9	41.1	21.7	2.9

ICH (12.9%) and 5–7 ICH metrics (55.9%) in a subsample (45–69 years, median age 55 years), was better than globally, mainly in high-income countries (0–2 ICH 32.2% and 5–7 ICH metrics 19.6%) [8]. It was also better than in rural South Africa (46% HIV positive, 38.7 years mean age) (7% had poor, 0–2 ideal metrics), and 53% 5–7 ICH metrics [11], and in Uganda (≥ 16 years, mean age 41.5 years in women and 36.8 years in men, 50.0% had 5–7 ICH metrics) [12]. The proportion of all seven ICH metrics (3.3%), intermediate ICH (≥ 1 metric in the intermediate category and none in the poor category) (15.3%) and poor ICH (≥ 1 metric in poor category) (81.5%) in this study, was similar to a study in Uganda ($N = 857$, ≥ 16 years, mean age 41.5 years in women and 36.8 years in men), (3.2% had 7 ICH metrics) [12], urban women in Ghana (mean age 45 years) 0.3% all seven ICH metrics, 25.0% in ≥ 1 metric in the intermediate category and none in the poor category 74.7% ≥ 1 metric in poor category [13], and in rural area in Northwest China all 7 ICH metrics (0.0%), intermediate (no poor health metrics and at least one intermediate) (18.0%), poor (any poor ICH metric) (82%) [16]. Our findings indicate that the prevalence of ICH is high but still efforts are needed to promote ICH to prevent CVD in Malawi.

Similar to the four best global estimates [10], this study showed that PA (97.8%), TC (95.0%), smoking (84.0%) and FBG (92.7%) had the highest prevalence of ideal status, while similar to the poorest global estimates [10], healthy diet (11.3%) had the poorest prevalence of ideal status in this study. The estimates of ideal PA (97.8%) in this study seem higher than global estimates of PA (40.6%) and ideal BMI (81.8%) are double of the global figures of ideal BMI (40.3%) [10]. In the 2009 STEPS national survey (24–64 years) in Malawi, a similar rate of ideal BMI (78.1%) was observed [24]. The high prevalence of ideal PA (97.8%) in this national study seems to be confirmed in the 2009 STEPS survey in Malawi (91.5% physically active) [25]. A low ideal healthy diet (fruit and vegetable consumption) (11.3% < 4.5 servings/day) was also found in the 2009 Malawi STEPS survey (2.5% < 5 servings/day) [26]. The proportion of poor smoking was 20.9% among men and 1.7% among women in this study, which is lower than the 2009 Malawi STEPS survey (25.9%

Table 4: Associations with meeting 5-7 ideal cardiovascular health metrics.

Variable	Crude OR (95% CI)	p-value	Adjusted OR (95% CI)	p-value
All				
Age in years				
18–29	1 (Reference)		1 (Reference)	
30–49	0.48 (0.45, 0.66)	<0.001	0.49 (0.35, 0.68)	<0.001
50–69	0.24 (0.17, 0.33)	<0.001	0.25 (0.17, 0.36)	<0.001
Sex				
Male	1 (Reference)		1 (Reference)	
Female	0.88 (0.72, 1.08)	0.219	0.98 (0.75, 1.27)	0.883
Education				
None	1 (Reference)		1 (Reference)	
Standard 1–4	1.27 (0.67, 2.40)	0.459	1.03 (0.69, 1.53)	0.901
Standard 5–8	1.91 (1.03, 3.55)	0.041	1.34 (0.88, 2.04)	0.174
Secondary or more	2.21 (1.20, 4.08)	0.011	1.13 (0.70, 1.84)	0.615
Employment status				
Nonpaid or unemployed	1 (Reference)		1 (Reference)	
Employed or student	1.01 (0.82, 1.24)	0.927	0.97 (0.75, 1.25)	0.808
Residence				
Rural	1 (Reference)		1 (Reference)	
Urban	0.58 (0.43, 0.79)	<0.001	0.56 (0.40, 0.78)	<0.001
Men				
Age in years				
18–29	1 (Reference)		1 (Reference)	
30–49	0.48 (0.29, 0.79)	0.004	0.49 (0.29, 0.82)	<0.001
50–69	0.19 (0.11, 0.32)	<0.001	0.20 (0.11, 0.36)	<0.001
Education				
None	1 (Reference)		1 (Reference)	
Standard 1–4	1.38 (0.96, 1.98)	0.085	0.83 (0.40, 1.69)	0.599
Standard 5–8	1.91 (1.32, 2.77)	<0.001	1.20 (0.60, 2.43)	0.605
Secondary or more	1.64 (1.11, 2.43)	0.013	1.23 (0.60, 2.50)	0.571
Employment status				
Nonpaid or unemployed	1 (Reference)		1 (Reference)	
Employed or student	1.11 (0.76, 1.62)	0.577	1.03 (0.67, 1.57)	0.904
Residence				
Rural	1 (Reference)		1 (Reference)	
Urban	1.10 (0.72, 1.66)	0.666	1.04 (0.67, 1.62)	0.855
Women				
Age in years				
18–29	1 (Reference)		1 (Reference)	
30–49	0.49 (0.36, 0.66)	<0.001	0.48 (0.35, 0.67)	<0.001
50–69	0.30 (0.21, 0.43)	<0.001	0.29 (0.19, 0.43)	<0.001

(contd.)

Variable	Crude OR (95% CI)	p-value	Adjusted OR (95% CI)	p-value
Education				
None	1 (Reference)		1 (Reference)	
Standard 1–4	1.49 (0.98, 2.27)	0.063	1.19 (0.75, 1.89)	0.464
Standard 5–8	1.96 (1.24, 3.10)	0.004	1.50 (0.89, 2.51)	0.125
Secondary or more	1.20 (0.69, 2.08)	0.522	0.94 (0.51, 1.74)	0.839
Employment status				
Nonpaid or unemployed	1 (Reference)		1 (Reference)	
Employed or student	0.83 (0.65, 1.07)	0.147	0.90 (0.68, 1.19)	0.456
Residence				
Rural	1 (Reference)		1 (Reference)	
Urban	0.40 (0.28, 0.56)	<0.001	0.40 (0.27, 0.59)	<0.001

OR = Odds Ratio; CI = Confidence Intervals.

among men and 2.9% among women) [25]. Poor BP (15.6%) was in this study (18–69 years) lower than in the 2009 Malawi STEPS survey (33.2%, hypertension, 25–64 years) [27]. Poor and intermediate FBG (7.4%) and poor and intermediate TC (4.9%) were similar to the 2009 survey (impaired and raised FBG 9.8% and 8.7% TC ≥ 5.0 mmol/L) [25, 28]. Similar to the 2009 STEPS survey, smoking and raised blood pressure were more frequent in men than in women, while obesity and raised TC occurred more often in women than in men [25].

Similar to a study in Northwest China [16], this study found that the proportion of having all four ideal health factors (36.1%) was significantly higher than those with all four ideal health behaviours (7.3%). Both, the proportion of ICH health factors and ICH health behaviours were higher in women than in men, while in a study in rural Uganda, ICH health factors were higher in men than in women and ICH health behaviours were higher in women than in men [12]. This result may pinpoint that the promotion of healthy behaviours should be emphasised to improve ICH [16]. A healthy diet (fruit and vegetable intake) was the least prevalent health metric (11.3%) in this study. Lack of affordability and availability of fruit and vegetables may be an influencing factor for the low intake of FV [11, 29]. Consequently, population strategies and interventions targeting persons with low fruit and vegetable consumption are urgently needed to improve ICH in Malawi.

Consistent with previous research [10, 12–16, 19], ICH was higher among younger age groups, rural residence, and those with higher education in unadjusted analysis.

The nonsignificant sex differences may be explained by on the one hand, women had a higher rate of ideal smoking and ideal BP than men, and on the other hand, men had a higher rate of ideal BMI, PA and TC than women.

To improve ICH in the Malawian, ICH behaviours should be improved through multidisciplinary interventions in living individuals, health educators, policy makers, and public health professionals [30]. Comprehensive interventions may target promotion of healthy diets, body weight control, smoking cessation, and screening and control of high levels of blood pressure and blood sugar [31]. Findings of the study may help in the NCD policy and plan of action in Malawi.

Most cases of hypertension and diabetes remain undiagnosed, untreated, or inadequately controlled. Our review demonstrated that there is an increased risk of hypertension and diabetes at a younger age and often in individuals with relatively low or normal BMI.

Given limited human and financial resources, innovative models of care are required to mitigate the growing burden of NCDs in Malawi. More broadly, we identified two innovative models of care integration across Malawi [20, 21]. The first model of care in southern Malawi is an integrated chronic care clinic that utilizes an HIV program as a platform for various chronic care screening and treatment (hypertension, diabetes, asthma, and epilepsy, regardless of HIV status) [21]. This model of care allows patients with chronic conditions or HIV to be screened and treated at a single facility during a single visit. The second model of care from central Malawi reported leveraging an HIV service platform for the screening and treatment of hypertension [20]. These studies documented that the integration of hypertension and diabetes screening into an HIV clinic is feasible despite various challenges including frequent stock out

and dispensing of NCD drugs, patient flow, workload, and issues related to data monitoring and evaluation [20, 21].

A recent NCD model of care review in sub-Saharan Africa also revealed that leveraging existing human resources, decentralization of NCD care to primary health facilities, task redistribution including to lay health cadres, designing patient-centered quality of care, and continued training and mentorship are key to successes of NCD treatment and control [24]. Although community screening and sensitization on the need for NCD care is critical, strategies are needed to ensure better linkage and retention into care [17].

The study strength is the large nationally representative sample and using standardized WHO STEPS methodology and measures. Study limitations include the cross-sectional design, the assessment of some variables was assessed by self-report. Furthermore, we included only one healthy diet component (fruit and vegetable consumption) and not all five components of AHA healthy diet. The participants included two-thirds women, but data were weighted for sex and age to the Malawian population.

Conclusion

The proportion of 5–7 ICH metrics was high in Malawian adults. Both high-risk and population-wide intervention programmes targeting older adults and urban residents should be implemented in aiding to improve ICH in Malawi.

Data Accessibility Statement

'The data for the current study are publicly available at the World Health Organization NCD Microdata Repository (URL: <https://extranet.who.int/ncdsmicrodata/index.php/catalog>).'

Abbreviations

AHA = American Heart Association; BP = Blood pressure; BMI = Body Mass Index; CVD = Cardiovascular disease; DALYs = Disability adjusted life years; FBG = Fasting blood glucose; FV = Fruit and Vegetables; ICH = Ideal Cardiovascular Health; NCD = non-communicable disease; PA = Physical activity; STEPS = *STEP-wise approach to Surveillance*; TC = *Total cholesterol*.

Acknowledgements

The data source, the World Health Organization NCD Microdata Repository (URL: <https://extranet.who.int/ncdsmicrodata/index.php/catalog>), is hereby acknowledged.

Competing Interests

The authors have no competing interests to declare.

Author Contribution

All authors fulfil the criteria for authorship. SP and KP conceived and designed the research, performed statistical analysis, drafted the manuscript and made critical revision of the manuscript for key intellectual content. All authors read and approved the final version of the manuscript and have agreed to authorship and order of authorship for this manuscript.

References

1. **World Health Organization (WHO)**. Cardiovascular diseases (CVDs), 2017. [https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds)) (accessed 11 November 2020).
2. **GBD 2019 Diseases and Injuries Collaborators**. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: A systematic analysis for the Global Burden of Disease Study 2019. *Lancet*. 17 Oct 2020; 396(10258): 1204–1222. DOI: [https://doi.org/10.1016/S0140-6736\(20\)30925-9](https://doi.org/10.1016/S0140-6736(20)30925-9).
3. **Lloyd-Jones DM, Hong Y, Labarthe D**, et al. Defining and setting national goals for cardiovascular health promotion and disease reduction: The American Heart Association's strategic Impact Goal through 2020 and beyond. *Circulation*. 2 February 2010; 121(4): 586–613. DOI: <https://doi.org/10.1161/CIRCULATIONAHA.109.192703>.
4. **Huffman MD, Capewell S, Ning H**, et al. Cardiovascular health behavior and health factor changes (1988–2008) and projections to 2020: Results from the National Health and Nutrition Examination Surveys. *Circulation*. 2012; 125: 2595–2602. DOI: <https://doi.org/10.1161/CIRCULATIONAHA.111.070722>

5. **Younus A, Aneni EC, Spatz ES**, et al. A Systematic Review of the Prevalence and Outcomes of Ideal Cardiovascular Health in US and Non-US Populations. *Mayo Clin Proc.* May 2016; 91(5): 649–70. DOI: <https://doi.org/10.1016/j.mayocp.2016.01.019>.
6. **World Health Organization.** Noncommunicable Diseases (NCD) Country Profiles, Malawi, 2018. https://www.who.int/nmh/countries/mwi_en.pdf?ua=1 (accessed 25 November 2020).
7. **Amberbir A, Lin SH, Berman J**, et al. Systematic Review of Hypertension and Diabetes Burden, Risk Factors, and Interventions for Prevention and Control in Malawi: The NCD BRITE Consortium. *Glob Heart.* June 2019; 14(2): 109–118. DOI: <https://doi.org/10.1016/j.gheart.2019.05.001>.
8. **Price AJ, Crampin AC, Amberbir A**, et al. Prevalence of obesity, hypertension, and diabetes, and cascade of care in sub-Saharan Africa: A cross-sectional, population-based study in rural and urban Malawi. *Lancet Diabetes Endocrinol.* March 2018; 6(3): 208–222. DOI: [https://doi.org/10.1016/S2213-8587\(17\)30432-1](https://doi.org/10.1016/S2213-8587(17)30432-1).
9. **Keates AK, Mocumbi AO, Ntsekhe M, Sliwa K, Stewart S.** Cardiovascular disease in Africa: Epidemiological profile and challenges. *Nat Rev Cardiol.* May 2017; 14(5): 273–293. DOI: <https://doi.org/10.1038/nrcardio.2017.19>.
10. **Peng Y, Cao S, Yao Z, Wang Z.** Prevalence of the cardiovascular health status in adults: A systematic review and meta-analysis. *Nutr Metab Cardiovasc Dis.* December 2018; 28(12): 1197–1207. DOI: <https://doi.org/10.1016/j.numecd.2018.08.002>.
11. **Ketelaar EJ, Vos AG, Godijk NG**, et al. Ideal Cardiovascular Health Index and Its Determinants in a Rural South African Population. *Glob Heart.* 25 November 2020; 15(1): 76. DOI: <https://doi.org/10.5334/gh.801>.
12. **Magodoro IM, Feng M, North CM**, et al. Female sex and cardiovascular disease risk in rural Uganda: a cross-sectional, population-based study. *BMC Cardiovasc Disord.* 25 April 2019; 19(1): 96. DOI: <https://doi.org/10.1186/s12872-019-1072-9>.
13. **van Nieuwenhuizen B, Zafarmand MH, Beune E**, et al. Ideal cardiovascular health among Ghanaian populations in three European countries and rural and urban Ghana: The RODAM study. *Intern Emerg Med.* September 2018; 13(6): 845–856. DOI: <https://doi.org/10.1007/s11739-018-1846-6>.
14. **Machado LBM, Silva BLS, Garcia AP**, et al. Ideal cardiovascular health score at the ELSA-Brasil baseline and its association with sociodemographic characteristics. *Int J Cardiol.* 1 March 2018; 254: 333–337. DOI: <https://doi.org/10.1016/j.ijcard.2017.12.037>.
15. **Ghimire U, Shrestha N, Gyawali B, Pradhan PMS, Mishra SR.** Prevalence of American Heart Association defined ideal cardiovascular health metrics in Nepal: Findings from a nationally representative cross-sectional study. *Int Health.* 1 July 2020; 12(4): 325–331. DOI: <https://doi.org/10.1093/inthealth/ihz088>.
16. **Zhao Y, Yan H, Yang R**, et al. Status of cardiovascular health among adults in a rural area of North-west China: Results from a cross-sectional study. *Medicine (Baltimore).* July 2016; 95(28): e4245. DOI: <https://doi.org/10.1097/MD.0000000000004245>.
17. **McKenzie JA, Younger NO, Tulloch-Reid MK**, et al. Ideal cardiovascular health in urban Jamaica: Prevalence estimates and relationship to community property value, household assets and educational attainment: A cross-sectional study. *BMJ Open.* 15 December 2020; 10(12): e040664. DOI: <https://doi.org/10.1136/bmjopen-2020-040664>.
18. **Ren J, Guo XL, Lu ZL**, et al. Ideal cardiovascular health status and its association with socioeconomic factors in Chinese adults in Shandong, China. *BMC Public Health.* 7 September 2016; 16(1): 942. DOI: <https://doi.org/10.1186/s12889-016-3632-6>.
19. **Benziger CP, Zavala-Loayza JA, Bernabe-Ortiz A**, et al. Low prevalence of ideal cardiovascular health in Peru. *Heart.* August 2018; 104(15): 1251–1256. DOI: <https://doi.org/10.1136/heartjnl-2017-312255>.
20. **Velasquez-Melendez G, Felisbino-Mendes MS, Matozinhos FP**, et al. Ideal cardiovascular health prevalence in the Brazilian population – National Health Survey (2013). *Rev Bras Epidemiol.* December 2015; 18 Suppl 2: 97–108. English, Portuguese. DOI: <https://doi.org/10.1590/1980-5497201500060009>.
21. **Ministry of Health.** Malawi National STEPwise Survey for Non-Communicable Diseases Risk Factors 2017 Report. file:///C:/Users/user/Downloads/2017%20STEPS%20Survey%20Report%20Malawi.pdf (accessed 10 January 2021).
22. **Armstrong T, Bull F.** Development of the World Health Organization Global Physical Activity Questionnaire (GPAQ). *J. Public Health.* 2006; 14: 66–70. DOI: <https://doi.org/10.1007/s10389-006-0024-x>

23. **Folsom AR, Yatsuya H, Nettleton JA**, et al. Community prevalence of ideal cardiovascular health, by the American Heart Association definition, and relationship with cardiovascular disease incidence. *J Am Coll Cardiol*. 2011; 57(16): 1690–6. DOI: <https://doi.org/10.1016/j.jacc.2010.11.041>
24. **Msyamboza KP, Kathyola D, Dzewela T**. Anthropometric measurements and prevalence of underweight, overweight and obesity in adult Malawians: Nationwide population based NCD STEPS survey. *Pan Afr Med J*. 24 July 2013; 15: 108. DOI: <https://doi.org/10.11604/pamj.2013.15.108.2622>.
25. **Msyamboza KP, Ngwira B, Dzewela T**, et al. The burden of selected chronic non-communicable diseases and their risk factors in Malawi: Nationwide STEPS survey. *PLoS One*. 2011; 6(5): e20316. DOI: <https://doi.org/10.1371/journal.pone.0020316>.
26. **World Health Organization (WHO)**. Malawi STEPS Survey 2009 Fact Sheet. https://www.who.int/ncds/surveillance/steps/2009_Malawi_FactSheet_EN.pdf?ua=1 (accessed 11 November 2020).
27. **Msyamboza KP, Kathyola D, Dzewela T, Bowie C**. The burden of hypertension and its risk factors in Malawi: Nationwide population-based STEPS survey. *Int Health*. December 2012; 4(4): 246–52. DOI: <https://doi.org/10.1016/j.inhe.2012.09.005>.
28. **Msyamboza KP, Mvula CJ, Kathyola D**. Prevalence and correlates of diabetes mellitus in Malawi: Population-based national NCD STEPS survey. *BMC Endocr Disord*. 12 May 2014; 14: 41. DOI: <https://doi.org/10.1186/1472-6823-14-41>
29. **Seron P, Irazola V, Rubinstein A**, et al. Ideal Cardiovascular Health in the southern cone of Latin America. *Public Health*. March 2018; 156: 132–139. DOI: <https://doi.org/10.1016/j.puhe.2017.12.017>.
30. **Moghaddam MM, Mohebi R, Hosseini F**, et al. Distribution of ideal cardiovascular health in a community-based cohort of Middle East population. *Ann Saudi Med*. March-April 2014; 34(2): 134–42. DOI: <https://doi.org/10.5144/0256-4947.2014.134>.
31. **Gyawali B, Mishra SR, Virani SS, Kallestrup P**. Low levels of ideal cardiovascular health in a semi-urban population of Western Nepal: A population-based, cross-sectional study. *Heart Asia*. 10 January 2019; 11(1): e011131. DOI: <https://doi.org/10.1136/heartasia-2018-011131>.

How to cite this article: Pengpid S and Peltzer P. Ideal Cardiovascular Health in a Nationally Representative Population-Based Sample of Adults in Malawi. *Global Heart*. 2021; 16(1): 24. DOI: <https://doi.org/10.5334/gh.986>

Submitted: 04 January 2021

Accepted: 26 March 2021

Published: 20 April 2021

Copyright: © 2021 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.



Global Heart is a peer-reviewed open access journal published by Ubiquity Press.

OPEN ACCESS