


## available at www.sciencedirect.com







## CORRIGENDUM

## Corrigendum to 'Gamma-Glutamyltransferase and mortality risk from heart disease and stroke in Japanese men and women: NIPPON DATA 90'

Akira Fujiyoshi <sup>a,\*</sup>, Katsuyuki Miura <sup>a</sup>, Atsushi Hozawa <sup>b</sup>, Yoshitaka Murakami <sup>c</sup>, Naoyuki Takashima <sup>a</sup>, Nagako Okuda <sup>a,g</sup>, Takashi Kadowaki <sup>a</sup>, Yoshikuni Kita <sup>a</sup>, Tomonori Okamura <sup>d</sup>, Yasuyuki Nakamura <sup>e</sup>, Takehito Hayakawa <sup>f</sup>, Akira Okayama <sup>g</sup>, Hirotsugu Ueshima <sup>a,h</sup>, for the NIPPON DATA80/90 Research Group

The authors of this paper would like to point out the following errors that appeared in the published article:

On page 29, under ''Results'' section, ''The 25th, 50th, 75th, and 90th percentile levels of GGT were 8, 12, 17, 26, and 52 U/L for women, and 15, 24, 41, 76, and 158 U/L for men.'' should read ''The 25th, 50th, 75th, and 90th

percentile levels of GGT were 11, 15, 22, and 37 U/L for women, and 19, 31, 58, and 111 U/L for men."

Also, in the footnote (a) of Table 1 (page 31), "Cut-off GGT values (U/L) for 25th, 50th, 75th, and 90th percentile were 8, 12, 17, 26 and 52 in women and 15, 24, 41, 76 and 158 in men, respectively." should read "Cut-off GGT values (U/L) for 25th, 50th, 75th, and 90th percentile were 11, 15, 22, and 37 in women and 19, 31, 58, and 111 in men, respectively."

The authors apologise for these errors.

E-mail address: afujiy@belle.shiga-med.ac.jp (A. Fujiyoshi).

<sup>&</sup>lt;sup>a</sup> Department of Health Science, Shiga University of Medical Science, Setatsukinowa-cho, Otsu, Shiga 520-2192, Japan

<sup>&</sup>lt;sup>b</sup> Division of Epidemiology, Department of Public Health and Forensic Medicine, Tohoku University School of Medicine, 2-1 Seiryomachi, Aoba-ku, Sendai 980-8575, Japan

<sup>&</sup>lt;sup>c</sup> Department of Medical Statistics, Shiga University of Medical Science, Setatsukinowa-cho, Otsu, Shiga 520-2192, Japan

<sup>&</sup>lt;sup>d</sup> Department of Preventive Cardiology, National Cardiovascular Center, 5-7-1 Fujishiro-dai, Suita, Osaka 565-8565, Japan

<sup>&</sup>lt;sup>e</sup> Cardiovascular Epidemiology, Kyoto Women's University, 35 Kitahiyoshi-cho, Imakumano, Higashiyama-ku, Kyoto 605-8501, Japan

<sup>&</sup>lt;sup>f</sup> Fukushima Medical University, Department of Hygiene & Preventive Medicine, 1 Hikariga-oka, Fukushima City, Fukushima Prefecture 960-1295, Japan

<sup>&</sup>lt;sup>g</sup> The First Institute for Health Promotion and Health Care, Japan Anti-Tuberculosis Association, 1-3-12 Suido-bashi Building, Misaki-cho, Chiyoda-ku, Tokyo 101-0061, Japan

<sup>&</sup>lt;sup>h</sup> Lifestyle-Related Disease Prevention Center, Shiga University of Medical Science, Setatsukinowa-cho, Otsu, Shiga 520-2192, Japan

DOI of original article: 10.1016/j.cvdpc.2010.03.003.

<sup>\*</sup> Corresponding author. Tel.: +81 77 548 2191; fax: +81 77 543